

AD-A147 440

(VINYLOXY)FLUOROCYCLOTRIPHOSPHAZENES(U) VERMONT UNIV  
BURLINGTON DEPT OF CHEMISTRY C W ALLEN ET AL  
25 OCT 84 TR 15 N00014 77 C 0605


1/8

UNCLASSIFIED

T/G 7/3

NI

END  
DATE  
12 MAR  
84



12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 15	2. GOVT ACCESSION NO. AD-A147448	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) (Vinyloxy)fluorocyclotriphosphazenes		5. TYPE OF REPORT & PERIOD COVERED Technical Report
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Christopher W. Allen and Randall P. Bright		8. CONTRACT OR GRANT NUMBER(s) N001477C-0605 Φ
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of Vermont Burlington, Vermont 05405		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS Department of the Navy Office of Naval Research Arlington, VA 22217		12. REPORT DATE 5/18/84
		13. NUMBER OF PAGES 10
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release and sale; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Accepted for publication in Inorganica Chimica Acta		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) cyclophosphazenes NMR enolate anions monomers		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The reactions of the lithium enolate of acetaldehyde, $\text{LiOCHCH}_3$ , with hexafluorocyclotriphosphazene, $\text{N}_3\text{P}_3\text{F}_6$ , lead to the series of (vinyloxy)fluorocyclotriphosphazene, $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n$ ( $n=2-5$ ). The $^1\text{H}$ NMR data shows that throughout the entire series, the substituent is bound through the oxygen end of the ambidentate enolate. The reaction shows a reluctance to go past the trisubstituted stage and the fluorine atom in $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n$ cannot be removed even under forcing conditions. Evidence from the $^1\text{H}$ , $^{19}\text{F}$ and $^{31}\text{P}$ NMR (continued on reverse)		

DTIC  
SELECTED  
NOV 09 1984

AD-A147 448

DTIC FILE COPY

DD FORM 1473  
1 JAN 73

EDITION OF 1 NOV 68 IS OBSOLETE  
G/N 0102-014-6001

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

84 11 05 100

cont'd

20. ABSTRACT (continued)

spectra shows that a non-geminal pathway is exclusively followed. Differences in the reaction patterns with enolate anions followed by  $\text{N}_3\text{P}_3\text{F}_6$  and  $\text{N}_3\text{P}_3\text{Cl}_6$  are discussed.

OFFICE OF NAVAL RESEARCH

Contract N001477C-0605

Project NR 356-663

Technical Report No. 15.

(Vinyloxy)fluorocyclotriphosphazenes

by

Christopher W. Allen and Randall P. Bright

Accepted for Publication in  
Inorganica Chimica Acta

University of Vermont  
Department of Chemistry  
Burlington, Vermont 05405

October 25, 1984

Accession For	
NTIS	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Accession Codes	
Distribution	
Dist	

Reproduction in whole or in part is permitted for any purposes  
of the United States Government.

This document has been approved for public release and sale;  
it's distribution is unlimited.

## (Vinylxy)fluorocyclotriphosphazenes.

Christopher W. Allen\* and Randall P. Bright  
Department of Chemistry, University of Vermont,  
Burlington, Vermont 05405, U.S.A.

Received

The reactions of the lithium enolate of acetaldehyde,  $\text{LiOCHCH}_2$ , with hexafluorocyclotriphosphazene,  $\text{N}_3\text{P}_3\text{F}_6$ , lead to the series of (vinylxy)fluorocyclotriphosphazene,  $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH=CH}_2)_n$  ( $n=2-5$ ). The  $^1\text{H}$  NMR data shows that throughout the entire series, the substituent is bound through the oxygen end of the ambidentate enolate. The reaction shows a reluctance to go past the trisubstituted stage and the fluorine atom in  $\text{N}_3\text{P}_3\text{F}(\text{OCH=CH}_2)_5$  can not be removed even under forcing conditions. Evidence from the  $^1\text{H}$ ,  $^{19}\text{F}$  and  $^{31}\text{P}$  NMR spectra shows that a non-geminal pathway is exclusively followed. Differences in the reaction patterns with enolate anions followed by  $\text{N}_3\text{P}_3\text{F}_6$  and  $\text{N}_3\text{P}_3\text{Cl}_6$  are discussed.

### Introduction

The reactions of enolate anions with halocyclotriphosphazenes occur exclusively at the oxygen end of the ambidentate nucleophile giving rise to (alkenoxo)phosphazenes [1]. One of these materials, the mono(vinylxy)penta-chlorotriphosphazene,  $\text{N}_3\text{P}_3\text{Cl}_5\text{OCH=CH}_2$ , has proven to be a valuable monomer for the synthesis of organofunctional phosphazene polymers [2]. Extensions of the enolate anion reaction have allowed for the preparation of a variety of derivatives of the type  $\text{N}_3\text{P}_3\text{Cl}_4(\text{X})\text{OCR=CH}_2$  ( $\text{X}=\text{Cl}$ ;  $\text{R}=\text{H}, \text{CH}_3, \text{C}_6\text{H}_5$ ;  $\text{X}=\text{CH}_3$ ,  $\text{R}=\text{H}, \text{CH}_3, \text{C}_6\text{H}_5$ ) [3]. The stereochemical course of the reaction of the enolate anion of acetaldehyde with  $\text{N}_3\text{P}_3\text{Cl}_6$  has been investigated [4]. All possible positional and stereoisomers in the series  $\text{N}_3\text{P}_3\text{Cl}_{6-n}(\text{OCH=CH}_2)_n$  ( $n=1-6$ ) have been detected

\*Author to whom correspondence should be addressed.

by  $^{31}\text{P}$  nmr spectroscopy. Although all the geminal isomers were detected, a predominantly non-geminal reaction pathway was observed. The non-geminal pattern is also observed in the reaction of aryloxides [5], alkoxides [6] and aryloxy steroid salts [7] with  $\text{N}_3\text{P}_3\text{Cl}_6$ . The corresponding fluorophosphazene reactions have received much less attention. Derivatives of  $\text{N}_3\text{P}_3\text{F}_6$  include  $\text{N}_3\text{P}_3\text{F}_5\text{OR}$  ( $\text{R}=\text{CH}_3, \text{C}_2\text{H}_5$ ) [8],  $\text{N}_3\text{P}_3\text{F}_5\text{OCR}=\text{CH}_2$  ( $\text{R}=\text{H}, \text{C}_6\text{H}_5$ ) [1],  $\text{N}_3\text{P}_3\text{F}_5\text{OC}(\text{CH}_2)_4\text{CH}_2$  [1] and 2,4,6- $\text{N}_3\text{P}_3\text{F}_3(\text{OCH}_2\text{C}_3\text{F}_7)_3$  [9]. The reactions of  $\text{N}_3\text{P}_3\text{X}_5\text{N}=\text{P}(\text{C}_6\text{H}_5)_3$  ( $\text{X}=\text{Cl}, \text{F}$ ) with the methoxide anion have recently been investigated [10]. In this paper, we present the synthesis of the series of (vinylxy)fluorocyclotriphosphazenes,  $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n$  ( $n=2-5$ ). The purpose for undertaking this investigation was to compare the reaction pathway followed by a specific oxygen base with  $\text{N}_3\text{P}_3\text{F}_6$  and  $\text{N}_3\text{P}_3\text{Cl}_6$  and to expand the range of available organofunctional phosphazenes.

### Experimental

Hexachlorocyclotriphosphazene (Firestone Corp.) was converted to hexafluorocyclotriphosphazene ( $\text{N}_3\text{P}_3\text{F}_6$ ) by a previously reported procedure [11]. n-Butyl lithium (1.6 M solution in hexane) was obtained from Aldrich. Tetrahydrofuran (THF) (Aldrich) was distilled from sodium-benzophenone ketyl. Hexanes and 35-55°C, benzene\*\* (Fisher) were distilled by standard procedures. NMR spectra (in  $\text{CDCl}_3$ ) were recorded on a Bruker WM250 spectrometer operating at 250.1 MHz ( $^1\text{H}$ ), 235.2 ( $^{19}\text{F}$ ) and 101.2 MHz ( $^{31}\text{P}$ ). Tetramethylsilane,  $\text{Me}_4\text{Si}$  (for  $^1\text{H}$  NMR) and fluorotrichloromethane,  $\text{CFCl}_3$  (for  $^{19}\text{F}$  NMR) were used as internal references. For  $^{31}\text{P}$  NMR, 85%  $\text{H}_3\text{PO}_4$  was used as an external standard. Chemical shifts upfield to the reference are assigned a negative sign.  $^{31}\text{P}$  NMR spectra were recorded under conditions of broad band decoupling. Mass spectra were determined on a Finnegan 4610 spectrometer operating at 80 eV. Elemental analyses were performed by Integral Microanalytical Laboratories.

\*\* Benzene is a suspected carcinogen, use only in a well-ventilated hood.

Preparation of  $N_3P_3F_4(OCH=CH_2)_2$ . A solution of 40 ml (0.064 mol) of n-butyllithium was added to 80 ml of tetrahydrofuran in an apparatus described elsewhere [12]. The resulting mixture was stirred for 16 hours at room temperature to allow for complete formation of the lithium enolate,  $LiOCH=CH_2$ , which was then added directly to 8.00g (0.032 mol) of  $N_3P_3F_6$  in 100 ml of THF at 0°C. The reaction mixture was heated to reflux. After removal of solvent, the resulting oil was distilled (bp 45-48°C at 0.10 mmHg) to yield 2.75 of crude product. A 2.00g sample of this material was subject to flash chromatography using benzene/hexanes (1/4) as an eluent. A 1.38 g (20.0% of theory) sample of a water-white liquid was obtained. Attempts at chromatographic separation of isomers failed. Anal. Calcd. for  $N_3P_3F_4O_2C_4H_6$ : C, 16.17; H, 2.02; mol. wt. 297. Found: C, 16.67; H, 1.94; mol. wt. 297 (mass spectrum).

Preparation of  $N_3P_3F_3(OCH=CH_2)_3$  and  $N_3P_3F_2(OCH=CH_2)_4$ . The lithium enolate, prepared from 100 ml of n-butyllithium (0.160 mol), was added to 10.0 g of  $N_3P_3F_6$  (0.04 mol) as described above. After removal of the solvent, the resulting oil was distilled (35-42°C at 0.02 mmHg) to give 4.65 g of crude material. A 3.00 g sample of this oil was subject to flash chromatography, as described above, resulting in the isolation of two products: 1.06 g (12.8% of theory) of a water-white liquid which was identified as  $N_3P_3F_3(OCH=CH_2)_3$  was obtained first. Anal. Calcd. for  $N_3P_3F_3O_3C_6H_9$ : C, 22.43; H, 2.80; mol. wt. 321. Found: C, 23.14; H, 2.68; mol. wt. 321 (mass spectrum). A second fraction contained 0.98 g (11.0% of theory) of a water-white liquid, which was identified as  $N_3P_3F_2(OCH=CH_2)_4$ , was also obtained. Anal. Calcd. for  $N_3P_3F_2O_4C_8H_{12}$ : C, 27.83; H, 3.48; mol. wt. 345. Found: C, 28.54; H, 3.52; mol. wt. 345 (mass spectrum). Attempts at chromatographic separation of isomers for the tris and tetrakis derivatives failed.

Preparation of  $N_3P_3F(OCH=CH_2)_5$ . The lithium enolate, prepared from 100



ml of n-butyllithium (0.16 mol), was added to 6.00g (0.024 mol) of  $N_3P_3F_6$  as described above. The oil remaining after removal of the solvent was distilled (bp 45–50°C at 0.02 mmHg) to give 0.94g of crude product. Chromatographic separation, as described above, gave primarily the tri and tetrasubstituted materials  $N_3P_3F_{6-n}(OCH=CH_2)_n$  ( $n=3,4$ ) along with 0.24g (2.7% of theory) of a water-white liquid which was identified as  $N_3P_3F(OCH=CH_2)_5$ . Anal. Calcd. for  $N_3P_3FO_5C_{10}H_{15}$ : mol. wt. 369. Found: Mol. wt. 369 (mass spectrum).

Attempted Preparation of  $N_3P_3(OCH=CH_2)_6$  from  $N_3P_3F_6$ . Several reactions were carried out with up to 10 molar equivalents of the enolate anion and reaction times of up to 120 hours. The major product obtained with these reactions was the tetrakis derivative along with small amounts of the tris and pentakis derivatives.

#### Results and Discussion

Previous studies have shown that attack on the phosphorus atom in formation of the monosubstituted derivatives of  $N_3P_3X_6$  ( $X=F[1], Cl[1,3]$ ) and the entire series of  $N_3P_3Cl_6[4]$  derivatives occurs at the oxygen end of the ambident enolate anion. These chemical precedents along with the close similarity of the  $^1H$  nmr spectra of the series  $N_3P_3F_{6-n}(OCH=CH_2)_n$  ( $n=2-5$ ) (Table I) to the previously reported [1] spectrum of  $N_3P_3F_5(OCH=CH_2)$  indicates that the new materials reported in this study are (vinyl oxy)fluorocyclotriphosphazenes.

The  $^{31}P$  nmr spectrum (Table II), of the products obtained at the stage of distribution,  $N_3P_3F_4(OCH=CH_2)_2$ , shows a large complex triplet, attributed to the  $^nPF_2$  center, and a complex doublet, corresponding to the  $^nPF(OCH=CH_2)$  centers. If a geminal isomer were present, a signal corresponding to a  $^nP(OCH=CH_2)_2$  center would be observed in the high field region of the spectrum, and would not possess a large, one bond phosphorus-fluorine coupling constant. No such signal was observed in the spectrum, thus, at the stage of distribution, the reaction proceeds exclusively by a non-geminal pathway. The  $^{19}F$  nmr spectrum

(Table II) shows two complex doublets corresponding to the  $\text{PF}_2$  and  $\text{PF}(\text{OCH}=\text{CH}_2)$  centers. The relative amounts of cis and trans isomers in  $\text{N}_3\text{P}_3\text{F}_4(\text{OCH}=\text{CH}_2)_2$  is unclear. The triplet in the  $^{31}\text{P}$  nmr spectrum is indicative of either a trans isomer or an equivalence of the  $^1J_{\text{PF}}$  coupling constants in the  $\text{PF}_2$  center since the two fluorine atoms in the  $\text{PF}_2$  center are inequivalent in the cis isomer. The Hx signals in the  $^1\text{H}$  nmr spectrum show a doubling of lines over what is observed in  $\text{N}_3\text{P}_3\text{F}_5\text{OCH}=\text{CH}_2$ . This observation combined with the fact that nearly equal amounts of cis and trans- $\text{N}_3\text{P}_3\text{Cl}_4(\text{OCH}=\text{CH}_2)_2$  are obtained in the corresponding reaction with  $\text{N}_3\text{P}_3\text{Cl}_6$  [4] suggests a cis/trans- $\text{N}_3\text{P}_3\text{F}_4(\text{OCH}=\text{CH}_2)_2$  mixture. The exclusive formation of non-geminal products in the  $\text{N}_3\text{P}_3\text{F}_6$  reaction is in contrast to the  $\text{N}_3\text{P}_3\text{Cl}_6$  reaction where a small amount of the geminal isomer is also observed [4].

In the trisubstituted derivative,  $\text{N}_3\text{P}_3\text{F}_3(\text{OCH}=\text{CH}_2)_3$ , the  $^{31}\text{P}$  NMR spectrum (Table II) shows a complex doublet, which can be assigned to the  $\text{PF}(\text{OCH}=\text{CH}_2)$  centers. No resonance is observed for either a  $\text{PF}_2$  or a  $\text{P}(\text{OCH}=\text{CH}_2)_2$  center, which would arise from a geminal (2,2,4-) substitution pattern. The  $^{19}\text{F}$  spectrum (Table II) allows for verification of this observation, and shows only a complex fluorine resonance in the  $\text{PF}(\text{OCH}=\text{CH}_2)$  region. For a geminal isomer to be present, a signal would be present in the  $\text{PF}_2$  region of the spectrum. Thus, at the stage of trisubstitution, only the formation of the non-geminal isomers is observed. This is in contrast to the observation of the formation of both the geminal and non-geminal isomers in the corresponding reaction of  $\text{N}_3\text{P}_3\text{Cl}_6$  leading to the trisubstituted derivative  $\text{N}_3\text{P}_3\text{Cl}_3(\text{OCH}=\text{CH}_2)_3$  [4]. The complexity of the  $^1\text{H}$ ,  $^{19}\text{F}$  and  $^{31}\text{P}$  spectra suggest the existence of both cis and trans- $\text{N}_3\text{P}_3\text{F}_3(\text{OCH}=\text{CH}_2)_3$ .

The  $^{31}\text{P}$  spectrum (Table II) of the tetrakis derivative,  $\text{N}_3\text{P}_3\text{F}_2(\text{OCH}=\text{CH}_2)_4$ , shows two resonances, a complex doublet and complex smaller multiplet. The doublet arises from the  $\text{PF}(\text{OCH}=\text{CH}_2)$  centers, while the small multiplet can be

assigned to the  $\text{P}(\text{OCH}=\text{CH}_2)_2$  center. If the geminal tetrakis isomer were present, a signal would occur which would correspond to a  $\text{PF}_2$  center. No such signal is observed. The  $^{19}\text{F}$  spectrum of the tetrakis derivative,  $\text{N}_3\text{P}_3\text{F}_2^-(\text{OCH}=\text{CH}_2)_4$  (Table II), shows a complex doublet in the  $\text{PF}(\text{OCH}=\text{CH}_2)$  range arising from the large phosphorus-fluorine coupling. A geminal isomer would show a signal corresponding to a  $\text{PF}_2$  center. Since no such signal is seen, the trend which is observed for the lower members of the series is followed at the stage of tetrakis substitution, i.e., non-geminal isomers are formed exclusively. The  $^1\text{H}$  nmr spectrum of the tetrakis derivative, shows extra signals which are due to the presence of a  $\text{P}(\text{OCH}=\text{CH}_2)_2$  center in the molecule.

The pentakis derivative,  $\text{N}_3\text{P}_3\text{F}(\text{OCH}=\text{CH}_2)_5$ , exhibits a large doublet and a small multiplet in the  $^{31}\text{P}$  nmr spectrum (Table II). Both resonances exhibit extensive coupling and second order effects. The doublet arises from the single  $\text{PF}(\text{OCH}=\text{CH}_2)$  resonance, while the singlet corresponds to the two  $\text{P}(\text{OCH}=\text{CH}_2)_2$  centers. The spectrum may be more appropriately be viewed as a second-order  $\text{AB}_2\text{X}$  system. The  $^{19}\text{F}$  NMR (Table II) consists of a doublet with extensive second order effects. The structure of the molecule is confirmed by the  $^1\text{H}$  NMR spectrum (Table VIII), which shows signals corresponding both to the  $\text{P}(\text{OCH}=\text{CH}_2)_2$  and the  $\text{PF}(\text{OCH}=\text{CH}_2)$  centers. Integration of these regions shows the relative abundance of 4:1 respectively, appropriate for the pentakis derivative.

It should be noted that all attempts to prepare the hexakis derivative,  $\text{N}_3\text{P}_3(\text{OCH}=\text{CH}_2)_6$ , were unsuccessful. Several reactions were carried out with a 10 molar equivalent of the enolate anion, and reaction times of up to 120 hours. The major product obtained with these reactions was the tetrakis derivative. Small amounts of tris and pentakis were also isolated.

The product distribution observed in the preparation of the vinylfluoro-cyclotriphosphazenes indicates that substitution occurs preferentially at a

$\text{PF}_2$  rather than a  $\text{PF}(\text{OCH}=\text{CH}_2)$  center. Evidence for this assertion is the complete absence of the geminal substitution pathway, the reluctance of the reaction to proceed past the trisubstituted stage and the inability to go past the pentasubstituted material. The decreasing reactivity at higher levels of substitution has been observed in other systems following a non-geminal pathway and has been ascribed to decreased electrophilicity of the phosphorus centers due to electron release to the ring from the substituent [13]. Non-geminal substitution is favored when the substitution is electron releasing relative to the halogen. The availability of the non-geminal pathway in the preparation of  $\text{N}_3\text{P}_3\text{Cl}_{6-n}(\text{OCH}=\text{CH}_2)_n$  [4] but not in  $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n$  can be related to the poor leaving group ability of the fluoride ion in phosphazene substitution reactions [14]. If the product distribution is kinetically controlled, then the low yield of geminal products in the chlorophosphazene reaction indicates a higher activation energy for the geminal pathway. The large energy associated with phosphorus-fluorine bond cleavage makes the formation of the geminal isomers energetically prohibitive in the fluorophosphazene reactions.

#### Acknowledgements

This work was supported, in part, by the Office of Naval Research. We thank Firestone Corporation for a generous gift of  $\text{N}_3\text{P}_3\text{Cl}_6$ .

### References

1. C.W. Allen, K. Ramachandran, R.P. Bright and J.C. Shaw, Inorg. Chim. Acta, 64, L109 (1982).
2. C.W. Allen, J. Polym. Sci., Polym. Symp., 70, 79 (1983).
3. P.J. Harris, M.A. Schwalke, V. Liu and B.L. Fisher, Inorg. Chem., 22, 1812 (1983).
4. K. Ramachandran and C.W. Allen, Inorg. Chem., 22, 1445 (1983).
5. C.T. Ford, F.E. Dickson and I.I. Bezman, Inorg. Chem., 4, 419 (1965); E.T. McSee, K. Okuhara and C.J. Morton, Ibid, 5, 450 (1966); D. Dell, B.W. Fitzsimmons, R. Keat and R.A. Shaw, J. Chem. Soc., A, 1966, 1680; W. Sulkowski, A.A. Volodin, K. Brandt, V.V. Kireev and V.V. Korshak, J. Gen. Chem. USSR (Engl. Transl.), 51, 1033 (1981).
6. J.L. Schmutz and H.R. Allcock, Inorg. Chem., 14, 2433 (1975); T.P. Zeleneva, I.V. Antonov and B.I. Stepanov, J. Gen. Chem. USSR (Engl. Transl.), 43, 1000 (1970).
7. H.R. Allcock, T.J. Fuller and K. Matsumura, J. Org. Chem., 46, 13 (1981).
8. E. Niecke, H. Thamm and O. Glemser, Z. Naturforsch., 26b, 366 (1971).
9. V.N. Prons, M.P. Grinblat, V.N. Sharov, A.L. Klebanski, J. Gen. Chem. USSR (Engl. Transl.), 47, 1149 (1977).
10. K.C. Kumaraswamy and S.S. Krishnamuthy, Phosphorus Sulfur, 18, 241 (1983).
11. T. Moeller, K. John and F. Tsang, Chem. Ind. (London), 1961, 347.
12. C.W. Allen, R.P. Bright, J.L. Desorcie, J.A. MacKay and K. Ramachandran, J. Chem. Educ., 57, 564 (1980).
13. J.M.E. Goldschmidt and E. Licht, J.C.S. Dalton, 1972, 728.
14. T. Moeller and S.G.J. Kokalis, J. Inorg. Nucl. Chem., 25, 1397 (1963).

TABLE I.  $^1\text{H}$  NMR Chemical Shift Data for  $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n$  ( $n=1-5$ ).<sup>a</sup>

Compound	$\delta_{\text{Ha}}^b$	$\delta_{\text{Hb}}^b$	$\delta_{\text{Hx}}^b$
$\text{N}_3\text{P}_3\text{F}_5\text{OCH}=\text{CH}_2^c$	5.13	4.85	6.48
$\text{N}_3\text{P}_3\text{F}_4(\text{OCH}=\text{CH}_2)_2$	5.09	4.80	6.48
$\text{N}_3\text{P}_3\text{F}_3(\text{OCH}=\text{CH}_2)_3$	5.06	4.77	6.49
$\text{N}_3\text{P}_3\text{F}_2(\text{OCH}=\text{CH}_2)_4$	5.06	4.73	6.51
	5.00	4.70	6.48
$\text{N}_3\text{P}_3\text{F}(\text{OCH}=\text{CH}_2)_5$	5.09	4.70	6.52
	4.97	4.65	6.48

a. Chemical shifts in ppm from TMS. b. Assignments are as follows:  
 $\text{=POC}=\text{C}-\text{Ha}$  c. Taken from reference 1.



Table II.  $^{31}\text{P}$  and  $^{19}\text{F}$  NMR Data for  $\text{N}_3\text{P}_3\text{F}_{6-n}(\text{OCH}=\text{CH}_2)_n^a$

Compound	$\equiv\text{PF}_2$ Centers			$\equiv\text{PF}(\text{OCH}=\text{CH}_2)$			$\equiv\text{P}(\text{OCH}=\text{CH}_2)_2$
	$\delta_{^{19}\text{F}}$	$\delta_{^{31}\text{P}}$	$^1J_{\text{PFB}}$	$\delta_{^{19}\text{F}}$	$\delta_{^{31}\text{P}}$	$^1J_{\text{PFB}}$	$\delta_{^{31}\text{P}}$
$\text{N}_3\text{P}_3\text{F}_5\text{OCH}=\text{CH}_2^c$	-69.5	11.0	910.0	-65.6	11.2	887.7	
$\text{N}_3\text{P}_3\text{F}_4(\text{OCH}=\text{CH}_2)_2$	-69.9	10.3	919.8	-66.1	10.6	895.8	
$\text{N}_3\text{P}_3\text{F}_3(\text{OCH}=\text{CH}_2)_3$				-65.7	11.0	871.5	
$\text{N}_3\text{P}_3\text{F}_2(\text{OCH}=\text{CH}_2)_4$				-65.5	12.3	903.0	11.1
$\text{N}_3\text{P}_3\text{F}(\text{OCH}=\text{CH}_2)_5$				-65.2	12.2	916.6	11.1

a. Chemical shifts in ppm; coupling constants in Hz. b. Approximate values from complex spectra.

c.  $^{31}\text{P}$  data taken from reference 1.

DL/413/83/01  
GEN/413-2

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. A. B. Amster Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		



ABSTRACTS DISTRIBUTION LIST, 356C

Professor A. G. MacDiarmid  
Department of Chemistry  
University of Pennsylvania  
Philadelphia, Pennsylvania 19174

Dr. E. Fischer, Code 2853  
Naval Ship Research and  
Development Center  
Annapolis, Maryland 21402

Professor H. Allcock  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Professor R. Lenz  
Department of Chemistry  
University of Massachusetts  
Amherst, Massachusetts 01002

Professor M. David Curtis  
Department of Chemistry  
University of Michigan  
Ann Arbor, Michigan 48105

Dr. J. Griffith  
Naval Research Laboratory  
Chemistry Section, Code 6120  
Washington, D.C. 20375

Professor G. Wnek  
Department of Materials Science  
and Engineering  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Mr. Samson Jennekke  
Honeywell Corporate Technology Center  
10701 Lyndale Avenue South  
Bloomington, Minnesota 55420

Dr. Richard M. Laine  
SRI International  
333 Ravenswood Avenue  
Menlo Park, California 94025

Dr. John McGrath  
Department of Chemical Engineering  
Virginia Polytechnic Institute  
Blacksburg, Virginia 24061

Dr. Adolf Amster  
Chemistry Division  
Naval Weapons Center  
China Lake, California 93555

Dr. William Tolles  
Code 6100  
Naval Research Laboratory  
Washington, D.C. 20375

Professor T. Katz  
Department of Chemistry  
Columbia University  
New York, New York 10027

Professor J. Salamone  
Department of Chemistry  
University of Lowell  
Lowell, Massachusetts 01854

Professor J. Chien  
Department of Chemistry  
University of Massachusetts  
Amherst, Massachusetts 01854

Professor William R. Krigbaum  
Department of Chemistry  
Duke University  
Durham, North Carolina 27706

Dr. R. Miller  
IBM Research Laboratory K42/282  
5600 Cottle Road  
San Jose, California 95193

DL/413/83/01  
356B/413-2

ABSTRACTS DISTRIBUTION LIST, 356B

Professor T. Marks  
Department of Chemistry  
Northwestern University  
Evanston, Illinois 60201

Professor Malcolm B. Polk  
Department of Chemistry  
Atlanta University  
Atlanta, Georgia 30314

Dr. Kurt Baum  
Fluorochem, Inc.  
680 S. Ayon Avenue  
Azusa, California 91702

Professor H. Ishida  
Department of Macromolecular Science  
Case Western University  
Cleveland, Ohio 44106

Professor Stephen Wellinohoff  
Department of Chemical Engineering  
University of Minnesota  
Minneapolis, Minnesota 55455

Professor G. Whitesides  
Department of Chemistry  
Harvard University  
Cambridge, Massachusetts 02138

Dr. K. Paciorek  
Ultrasystems, Inc.  
P.O. Box 19605  
Irvine, California 92715

Professor H. Hall  
Department of Chemistry  
University of Arizona  
Tucson, Arizona 85721

Professor D. Seyferth  
Department of Chemistry  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139